

POTENCY OF HONEY AS ANTIBACTERIAL AGENT AGAINST MULTIPLE ANTIBIOTIC RESISTANT PATHOGENS EVALUATED BY DIFFERENT METHODS

¹Mohamed Mustafa Aween, ¹Zaiton Hassan, ¹Nur Huda-Faujani,
¹Mohamed Muftah Emdakim and ²Belal Jamal Muhiaddin

¹Faculty of Science and Technology, Universiti Sains Islam Malaysia (USIM),
71800, Bandar Baru Nilai, Negeri Sembilan, Malaysia

²Faculty of Food Science and Technology, Universiti Putra Malaysia (UPM),
UPM Serdang, 43400, Selangor, Malaysia

Received 2014-08-04; Revised 2014-10-15; Accepted 2014-10-20

ABSTRACT

Honey is rich with complex natural components which could be useful as antibacterial agents or as preservative. Honey contains high concentration of sugars, low amount of water, high osmolality and often dark colour which influence its antibacterial activity. Disc diffusion, well method, micro dilution assay are methods commonly used to determine the antibacterial activity of honey. In this study, microtiter and microbial plate count were included to ascertain the potency of honey as antibacterial agent against multiple antibiotic resistant pathogenic bacteria (*Staphylococcus aureus*, *Salmonella Typhimurium*, *Escherichia coli*, *Bacillus subtilis* and *Pseudomonas aeruginosa*) with concentration of 0.2 g mL⁻¹. Results found that well diffusion method tends to give higher inhibitory zone than disc diffusion method but there was no correlation among the bacteria was observed except for *S. Typhimurium*, *E. coli* (R = 0.310, 0.505 and 0.316, respectively). Nan photometer assay and microtiter plates assay showed comparable results with moderately strong correlation (R² = 0.681 and 0.767, respectively) for *S. aureus* and *S. typhimurium*, but poor correlation was found for *E. coli*, *B. subtilis* and *P. aeruginosa* (R² = 0.441, 0.308 and 0.383, respectively). Determining the number of survivors by plating on agar after nanophotometer assay or microtiter plate assay had confirmed the effectiveness of honey as antimicrobial agent against target bacteria; which confirmed that honey has the potency to inhibit pathogens even at low concentration.

Keywords: Honey, Antibacterial Activity Assay, Multiple Antibiotic Resistant (MAR), Pathogenic Bacteria

1. INTRODUCTION

Honey not only a delicious food but are also known for its health and healing properties. Such concept is referred to in Surat El-Nahl, the Holy Quraan (verse 68 and 69). Honey has the ability to inhibit microorganisms including bacteria, spoilage fungi and yeast and viruses. The antibacterial effect of honey especially against Gram-positive bacteria are well documented (Molan,

1997; Bogdanov, 1997; Rozaini *et al.*, 2004; Tan *et al.*, 2009). Both bacteriostatic and bactericidal effects of honey have been reported for many strains especially pathogenic bacteria (Bogdanov *et al.*, 2008). It is suggested that Manuka honey (6% v/v) can be used against *Burkholderia cepacia* which causes pulmonary infections and chronic granulomatous disease in urinary tract infections and wound infections in hospitalized patients (Cooper *et al.*, 2000a; 2000b).

Corresponding Author: Mohamed Mustafa Aween, Faculty of Science and Technology, Universiti Sains Islam Malaysia (USIM), 71800, Bandar Baru Nilai, Negeri Sembilan, Malaysia

Honey has significant amount of glucose oxidase, catalase, ascorbic acid, flavonoids, phenolic acids, carotenoid derivatives, organic acids, Maillard reaction products, amino acids and proteins that contribute to its functional properties (Bogdanov *et al.*, 2008; Perez *et al.*, 2007). Honey contains cinnamic acid, antioxidant agent and some flavonoids which have been approved for antibacterial applications (Rahman *et al.*, 2010). Mohammed (2010) reported that Malaysian honey have antioxidant properties. The antioxidants compounds in honey may play a positive role in food safety beyond food preservation (Taormina *et al.*, 2001).

In Malaysia, research on honey has focused on different aspects. Tualang honey was reported to be effective for wound healing and wound burn management (Nasir *et al.*, 2010) and the effects of different types of honey on tensile strength evaluation of burn wound tissue healing were evaluated by Rozaini *et al.* (2004). The types of phenolic acids in Malaysian honey were suggested to be responsible for the antibacterial properties (Aljadi and Yusoff, 2003) while other local Malaysian honey showed antimicrobial activity on some human pathogens (Tumin *et al.*, 2005; Hassanain *et al.*, 2010; Zainol *et al.*, 2013). Propolis of Malaysian honey inhibited the growth of *Staphylococcus aureus* and *Escherichia coli* as reported by Rahman *et al.* (2010). Honey of Malaysia, Libya, New Zealand and Saudi Arabia contain lactic acid bacteria with antibacterial activity against selected Multiple Antibiotic Resistant (MAR) Gram-positive bacteria and Gram-negative bacteria (Aween *et al.*, 2012a; 2012b).

Two methods that are commonly used to evaluate the antibacterial activity are disc diffusion method (Bauer *et al.*, 1966) and well diffusion (Perez *et al.*, 1990). Since honey is a complex of substances, these methods may not reflect the potency of honey as antibacterial agent. Therefore, other methods namely nanophotometer assay, microtiter plates and microbial plate count methods were included in this study in an attempt to evaluate the effectiveness of the methods for determination of the antibacterial activity of honey on multiple antibiotic resistant pathogenic bacteria.

2. MATERIALS AND METHODS

2.1. Honey Samples

Nine samples of honey were obtained from different sources and coded as follows: Tualang honey (H026) and Acacia honey (H030, H031 and H032) from Malaysia, Al-Seder (H025), Kharoob (H028) and Hannon Honey (H020) from Libya and Manuka Honey 5+ (H027) and 10+ (H035)

from New Zealand. All samples were kept in glass bottles and stored at room temperature before experiment.

2.2. Honey Samples Preparation

Amount of honey used was based on the lowest concentration that showed antibacterial activity. Honey samples (0.2 g mL⁻¹ deionized water) were prepared and all honey samples were heated at 70°C for 10 min using water bath and filtered using 0.45 µL filters and then kept at 4°C for further study. Heating removed H₂O₂ and destroyed any contaminating microorganisms that may be present in honey samples.

2.3. Cultures of Pathogenic Bacteria

The pathogenic bacteria used in this study were *S. aureus*, *S. Typhimurium*, *E. coli*, *B. subtilis* and *P. Aeruginosa* obtained from Microbiology Laboratory Faculty of Science and Technology, Universiti Sains Islam Malaysia (USIM), Malaysia. The bacteria were grown on nutrient agar (Oxoid, UK) and nutrient broth (Oxoid, UK) at 37°C for 24 h and then kept at 4°C before further experiments.

2.4. Antibiotic Resistant Test of Target Bacteria

The target bacteria were tested for their resistant to antibiotics using disc diffusion method as described by Bauer *et al.* (1966). The antibiotics used were vancomycin (5 µm), cephalothin (30 µm), nalidixic acid (30 µm), Gentamycin (10 µm), streptomycin (10 µm), tetracycline (30 µm), bacitracine (10 µm), penicillin G (10 µm), chloramphenicol (30 µm) and polymyxin B (300 µm) (Sigma). The selection of antibiotics in this study was based on the common antibiotics used in medical practice and health therapy. Mean, standard deviation and MAR index were calculated for all target bacteria (Subramani and Vignesh, 2012).

2.5. Antibacterial Activity of Honey Samples using Disc Diffusion Method

Antibacterial activity of selected honey samples were determined by disc diffusion method on Nutrient Agar (NA) agar (Oxoid, UK) (Bauer *et al.*, 1966). The pathogenic bacteria cultures were swabbed on NA agar plates. Discs were overlaid with tested honey samples overnight and then dried at 45°C for 24 h using drier oven (BINDER, Germany). Discs were placed on swabbed agars and incubated at 37°C for 24 h. Inhibition zone diameter was carefully measured and the results were expressed in millimetre. This experiment was done in duplicate and mean with standard deviation were calculated.

2.6. Antibacterial Activity of Honey Samples using well Diffusion Method

The well diffusion method for antibacterial activity of honey was determined following the method of Perez *et al.* (1990) with slight modifications. Honey samples were prepared with concentration of 0.2 g mL⁻¹ (w/v) using deionized water. Overnight culture of pathogenic bacteria in nutrient broth (Oxoid, UK) was prepared. Nutrient agar was prepared and once its temperature reached 40°C, 1% of pathogenic bacteria (10⁹ CFU/ml) was added and mixed carefully. Amount of 25 mL of the nutrient agar with 1% of pathogenic bacteria was poured to petri dish plates and left under a laminar flow until the plates dried. Wells of 8 mm diameter was made using a sterile cork-borer with 8 mm diameter and the base of the wells were covered with nutrient agar (Oxoid, UK) and left to dry at room temperature. After that, 200 µL of prepared honey samples were poured to the wells individually and kept at 37°C for 24 h. The results were expressed by measuring the zones around the wells after diminution the well size. The experiment was done in duplicate and mean with standard deviation were calculated.

2.7. Antibacterial Activity of Honey Samples using Nanophotometer Assay

In Nanophotometer method, concentration of 0.2 g mL⁻¹ honey samples were prepared using deionized water. Pathogenic bacteria were inoculated to Nutrient broth (Oxoid, UK) and kept at 37°C. 1 mL of each honey sample was poured to 1 mL of pathogenic bacteria (10⁶ CFU/ml) in micro-titer plates and kept at 37°C for 24 h. 1 mL of each honey sample with 1 mL of nutrient broth without pathogenic bacteria was used as negative control and 1 mL of broth with pathogens with 1 mL of broth was used as positive control. The reading was determined as bacteria cells using Nanophotometer (IMPLEN, Germany) at wavelength of 600 nm. The results then were interpreted using the following formula:

Percentage of inhibition = [(Positive control absorbance-Sample absorbance)/Positive Control absorbance] × 100

2.8. Antibacterial Activity of Honey Samples using Microtiter Plates

Each honey sample was tested against the selected pathogenic bacteria in microtiter plate assay, following the method of Magnusson and Schnurer (2001) with some modifications. 100 µL of nutrient broth (NB)

containing 10⁶ pathogenic bacteria/ml were placed in the 96 wells plate and 150 µL honey samples (0.2 g mL⁻¹) were poured into the wells. The plates were incubated at 37°C for 24 h. Optical density of bacterial growth was measured at 630 nm using Elisa plate reader (BIOTEK, USA). Honey with Nutrient Broth (NB) without bacteria was used as negative control and Nutrient Broth (NB) with pathogenic bacteria was used as positive control. The results then were interpreted using the following formula:

Percentage of inhibition = (+ Control absorbance-Sample absorbance)/ + Control absorbance) × 100

2.9. Antibacterial Activity of Honey Samples using Total Plate Count Assay

Antibacterial activity of selected honey samples were determined by plate count method on Nutrient Agar (NA) (OXOID) following Aween *et al.* (2012b) with some modifications. 100 µL of overnight pathogenic bacteria culture in NB were added to 100 µL of honey sample and kept at 37°C for 24 h. After that, 10 µL of the mixture were spread on NA and incubated at 37°C for 24 h. Pathogenic bacteria in NB was used as positive control and honey sample with NB as negative control. The results were taken by counting the number of pathogenic bacteria colonies on NA plates after 24 h of incubation and expressed as log₁₀ CFU/ml. The enumeration of bacteria on plates followed standard microbiological procedure.

2.10. Statistical Analysis

All data were analyzed using Minitab 16 system to calculate the mean, standard deviation and percentage of inhibition. The correlation was carried out between the methods (disk diffusion method versus well diffusion method and microtiter plates versus nanophotometer) and the bacteria tested were determined (R = <0 no correlation, R = 0-0.7 poor correlation and R = 0.7-1 strong correlation).

3. RESULTS

3.1. Antibiotic Resistant of Selected Target Bacteria

The target bacteria showed high resistance to several antibiotics tested. The diameter of inhibition zone varies between 0 and 29 mm (Table 1). All target bacteria were not inhibited by Bacitracin, Polymyxin B, Penicillin G, Vancomycin and Streptomycin. However, *S. aureus* were totally resistant to polymyxin B, while *S.*

Typhimurium and *B. subtilis* were resistant to bacitracin, tetracycline, penicillin G, vancomycin, naladixic acid and Chloramphenicol. *E. coli* were resistant to Bacitracin, Gentamycin, Penicillin G and Vancomycin. Bacitracin, Tetracycline, Penicillin G, Vancomycin and Chloramphenicol were not effective against *P. aeruginosa*. MAR index was from 11 to 88%, *S. Typhimurium* and *B. subtilis* showed the highest MAR index percentage (88%) compared to *E. coli* (55%), *P. aeruginosa* (66%) and *S. aureus* (11%).

3.2. Antibacterial Activity of Honey Samples using Disc Diffusion Method

All the nine honey samples showed variable inhibitory activities against the target bacteria by the disc diffusion method (Table 2). The inhibitory

activity was significantly ($p < 0.5$) affected by the type of bacteria used but not with honey samples. *P. aeruginosa* was greatly inhibited by all honey samples as shown by inhibitory zone greater than 10.5 ± 4.94 mm except H032 which was less than (7.5 ± 0.70 mm). While *E. coli* and *S. Typhimurium* were inhibited but to a lesser extent (Table 2). Tualang honey (H026) from Malaysia showed the highest inhibitory activity against *S. aureus*, *S. Typhimurium* and *B. subtilis* (20.00 ± 1.41 , 14.00 ± 4.24 and 17.00 ± 0.00 mm respectively), Acacia honey (H031) from Malaysia showed the highest inhibitory zone of 18.50 ± 3.53 and 17.00 ± 0.00 mm against *E. coli* and *B. subtilis*, while against *P. aeruginosa* Al-Seder honey (H025) from Libya showed the highest activity.

Table 1. The antibacterial activities of selected antibiotics against target bacteria^a

Antibiotics	Target bacteria				
	<i>S. aureus</i>	<i>S. Typhimurium</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>
Bacitracin (10 µg)	6.50±0.70	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Gentamycin (120 µg)	11.00±0.00	14.50±0.70	0.00±0.00	13.00±0.00	13.00±0.00
Tetracycline (10 µg)	13.50±2.12	0.00±0.00	8.00±5.65	0.00±0.00	2.50±2.12
Naladixic acid (30 µg)	8.50±0.70	0.00±0.00	18.00±0.00	0.00±0.00	17.00±0.00
Polymyxin B (300 µg)	0.00±0.00	4.00±0.00	4.00±0.00	4.00±0.00	4.00±0.00
Penicillin G (5 µg)	29.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Vancomycin (5µm)	6.00±0.00	0.00±0.70	0.00±0.00	0.00±0.00	0.00±0.00
Streptomycin (10 µm)	6.00±0.00	3.00±0.00	8.50±0.70	3.00±0.00	7.50±0.70
Chloramphenicol (30 µg)	15.00±2.82	0.00±0.00	18.50±4.94	0.00±0.00	0.00±0.00
MAR index %	11	88	55	88	66

^aDiameter of inhibition zone around the discs (mm)

Table 2. Growth inhibition zone of target bacteria by honey samples by disc diffusion method^a

Honey sample	Target bacteria				
	<i>S. aureus</i>	<i>S. Typhimurium</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>
H020	16.50±0.70	11.50±4.94	13.00±4.24	15.50±0.70	15.00±0.00
H025	19.50±0.70	11.50±3.53	10.50±2.12	15.50±0.70	17.00±0.00
H026	20.00±1.41	14.00±4.24	13.50±3.53	17.00±0.00	16.00±0.00
H027	17.00±0.00	11.50±2.12	11.00±1.41	15.50±0.70	16.00±0.00
H028	17.00±1.41	10.50±3.53	11.50±3.53	14.00±0.00	15.00±0.00
H030	19.00±0.00	13.50±2.12	15.00±2.82	15.50±0.70	14.50±3.53
H031	16.50±3.53	12.00±1.41	18.50±3.53	17.00±0.00	12.50±4.94
H032	17.00±1.41	11.00±1.41	13.50±2.12	16.50±0.70	7.50±0.70
H035	18.00±1.41	11.50±3.53	14.50±2.12	15.50±0.70	10.50±4.94

^aDiameter of growth inhibitory zone was measured in mm after 24 h incubation at 37°C

3.3. Antibacterial Activity of Honey Samples using well Diffusion Method

In well diffusion method, all honey samples from different sources showed inhibitory activity against all the target pathogenic bacteria; however, in the current study the degree of inhibition was affected by type of bacteria; the growth inhibitory zone varies between 15.5 and 27.5 mm (Table 3 and Fig. 1). Among the bacteria evaluated, *S. aureus* was easily inhibited by all the tested honey samples, while *E. coli* was the most difficult to be inhibited. Growth of *S. aureus* was easily inhibited by Hannon honey, Libya (H020), Acacia honey, Malaysia (H031), Acacia honey, Malaysia (H032) and Manuka honey, New Zealand (H027) with inhibitory zone of 27.50, 25.00, 25.00 and 25.50 mm, respectively (Table 3 and Fig. 1). Growth of *S. Typhimurium* and *B. subtilis* were moderately inhibited by all honey samples with inhibitory zone between 15.50 to 19.50 mm. All honey samples also inhibited *P. aeruginosa* (16.00 to 18.50 mm inhibitory zone). Libyan honey Hannon honey (H020) showed the highest inhibitory activity

against all tested pathogenic bacteria (18.50 to 27.50) except against *E. coli* (15.00 mm), while Acacia honey (H032) was effective against *S. aureus* (25.00 mm), *S. Typhimurium* (18.00 mm) and *E. coli* (17.00 mm).

3.4. Antibacterial Activity of Honey Samples using Nanophotometer Assay

In Nanophotometer assay the survival of target bacteria was determined using nanophotometer and the results were expressed as percentage inhibition from survival (CFU/ml) of the bacteria after treatment with honey (Table 4). The percentage of inhibition of all honey samples ranged from 13.17 to 100%. Al-Seder honey (H025) showed total inhibition (100%) against *S. aureus*, *E. coli* and *P. aeruginosa*; Acacia honey (H030) showed inhibitory activity against *S. aureus* and *S. Typhimurium* (100 and 99.09%). *S. aureus* and *E. coli* were easily inhibited by all tested honey samples with percentage of 94.26 to 100 and 83.36 to 100, respectively. All samples of honey showed inhibitory activity against *P. aeruginosa* except H031 (13.17%).

Table 3. Growth inhibition zone of target bacteria by honey samples using well method^a

Honey sample	Target bacteria				
	<i>S. aureus</i>	<i>S. Typhimurium</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>
H020	27.50±0.70	19.00±1.41	15.00±2.82	19.50±0.70	18.50±2.12
H025	22.50±0.70	16.50±0.70	15.50±0.70	17.50±0.70	16.00±0.00
H026	24.50±0.70	18.00±1.41	16.50±0.70	17.50±0.70	16.00±0.00
H027	25.50±2.12	17.00±2.12	14.00±0.00	19.00±0.00	17.50±0.70
H028	24.50±3.53	15.50±0.70	14.00±0.00	18.00±0.00	16.50±0.70
H030	23.50±0.70	17.50±0.70	15.00±0.00	18.50±0.70	16.00±0.00
H031	25.00±0.70	17.50±0.70	16.50±0.70	17.50±0.70	16.00±0.00
H032	25.00±0.00	18.00±0.00	17.00±0.00	18.00±1.41	16.00±0.00
H035	24.00±0.00	18.50±0.70	15.00±0.70	17.50±0.70	16.00±0.00

^aDiameter of growth inhibitory zone was measured in mm after 24 h incubation at 37°C, ± = SD

Table 4. Percentage of inhibition of target bacteria by honey samples using Nanophotometer assay^a

Honey sample	Target bacteria (% inhibition)				
	<i>S. aureus</i>	<i>S. Typhimurium</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>
H020	94.26	78.47	83.36	82.18	93.92
H025	100.00	83.68	100.00	92.78	100.00
H026	100.00	93.59	99.82	97.81	99.25
H027	99.15	82.28	90.03	84.67	96.75
H028	99.46	87.18	99.17	87.36	97.54
H030	100.00	99.09	97.18	96.06	96.93
H031	97.92	79.57	94.84	95.82	13.17
H032	98.40	88.08	96.92	95.82	93.27
H035	99.15	94.09	92.11	90.64	70.40

^aPercentage of inhibition = [(+ Control absorbance - Sample absorbance)/+Control absorbance] × 100




Fig. 1. Growth inhibition zone of honey samples against pathogenic bacteria by well method at 37°C after 24 h of incubation

Table 5. Percentage of inhibition of target bacteria by honey samples using microtiter plates

Honey sample	Target bacteria (% inhibition)				
	<i>S. aureus</i>	<i>S. Typhimurium</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>
H020	70.14	74.63	70.93	74.44	64.04
H025	98.48	86.52	85.43	89.17	93.73
H026	100.00	92.38	90.69	95.22	89.08
H027	99.33	89.22	96.89	96.25	97.49
H028	80.37	85.06	83.28	85.17	74.50
H030	89.00	92.32	88.47	91.02	80.05
H031	70.61	77.15	76.12	74.14	73.25
H032	98.48	93.79	93.54	95.58	87.56
H035	97.53	89.45	82.52	86.40	75.76

^aPercentage of inhibition= [(+ Control absorbance -Sample absorbance)/+Control absorbance] ×100

3.5. Antibacterial Activity of Honey Samples using Microtiter Plates

Microtiter plates measures growth of bacteria by turbidity at OD₆₃₀. All tested honey samples exhibited high inhibitory activity against all target pathogenic bacteria. Percentage of inhibition of target bacteria

ranged from 60 to 100% by all honey samples within 24 h of incubation (Table 5). The growth of *S. aureus* was totally inhibited (100%) by Tualang honey (H026), while Acacia honey (H032) showed the highest inhibitory activity against *S. Typhimurium* (93.79%). The lowest activity was obtained from Hannon honey (H020) against all tested pathogens (64.04-74.63%).

Table 6. Growth inhibition of target bacteria by honey samples using total plate count method^a

Honey sample	Target bacteria (CFU/ml)				
	<i>S. aureus</i>	<i>S. Typhimurium</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>
H020	<10 (est.)	10 ⁵	<10 (est.)	10 ⁵	10 ⁵
H025	10 ⁵	10 ⁵	<10 (est.)	10 ⁵	<10 (est.)
H026	<10 (est.)	<10 (est.)	<10 (est.)	<10 (est.)	<10 (est.)
H027	10 ⁵	10 ⁵	<10 (est.)	10 ⁵	10 ⁵
H028	<10 (est.)	10 ⁵	<10 (est.)	<10 (est.)	<10 (est.)
H030	<10 (est.)	<10 (est.)	<10 (est.)	<10 (est.)	10 ³
H031	10 ³	10 ³	10 ⁵	10 ³	10 ⁵
H032	<10 (est.)	<10 (est.)	<10 (est.)	<10 (est.)	10 ⁵
H035	<10 (est.)	10 ³	10 ⁵	10 ⁵	10 ⁵
+ Control	10 ¹⁶	10 ¹⁵	10 ¹⁵	10 ¹⁵	10 ¹⁶

^a10: Number of colonies

3.6. Antibacterial Activity of Honey Samples using total Plate Count Assay

Survival of target bacteria as evaluated by plate count method was significantly affected by honey samples and by type of bacteria (Table 6). Tualang Honey (H026) had inhibited all tested pathogenic bacteria, followed by H030 and Acacia honey (H032) which inhibited all tested bacteria except *P. aeruginosa*. Growth of *E. coli* was totally inhibited by all honey samples except Acacia honey (H031) and Manuka honey (H035). *S. aureus* was not sensitive to Al-Seder honey (H025) and Manuka honey (H027), but inhibited by all the others. Hannon honey (H020) and Al-Seder honey (H025) were not active against *S. Typhimurium* and *B. subtilis*. Generally, all the honey samples showed good antibacterial activity against tested pathogenic bacteria compared to the positive control as evaluated by this method.

4. DISCUSSION

The occurrence of MAR bacterial strains is a public health concerns due to the bacteria are not easily killed by common antibiotics that normally used for health therapy. Some strains of *Staphylococcus* species were resistant to several antibiotics (Salvatore *et al.*, 2010); *S. aureus* was resistant to ampicillin, kanamycin and oxytetracycline. The *S. aureus* used in this study demonstrated low MAR index 11% compared *E. coli* and *P. aeruginosa* with 55 and 66%, respectively. *S. Typhimurium* and *B. subtilis* showed highest MAR index (88%) compared to other tested bacteria. Similarly, multiple antibiotic resistant bacteria was shown by several tested pathogens including *S. Typhimurium* ATCC13311 that was resistant to 10 antibiotics especially to bacitracin, cephalothin, penicillin G, vancomycin and streptomycin; *E. coli* ATCC25922 was

resistant to bacitracin, penicillin G and vancomycin observed by Aween *et al.* (2012b).

The presence of multiple resistance pathogenic bacteria has led to the investigation of natural effective alternatives to common antibiotics used in medical practice and health therapy. The results of presence study showed that honey available in Malaysia can be used as antibacterial agent to prevent and control infections which are caused by the pathogenic bacteria. Five different methods were used to evaluate the antibacterial activity of different honey samples and the results showed no correlation ($R = < 0$) between disc diffusion method and well diffusion method except for *S. Typhimurium*, *E. coli* and *P. aeruginosa* which was poor ($R = 0.310, 0.505$ and 0.316 , respectively). Well diffusion method was more suitable method for detecting the susceptibility of bacteria to antibacterial substances compared to disc diffusion. In contrast, the disc diffusion method is mainly used as a qualitative test for detecting the susceptibility of bacteria to antimicrobial substances (Mandal and Mandal, 2011). Disc diffusion is based on the ability of molecules to diffuse into the agar; while well method, nanophotometer assay and microtiter plates allowed direct contact of the bacterial surfaces to the compounds. It was also observed that there was correlation ($R = 0.308$ to 0.767) between nanophotometer assay and microtiter plates method.

Earlier reports showed that Egyptian clover honey tested for its antibacterial effect against antibiotic resistant strains of *E. coli* and *S. Typhimurium* using disc diffusion method was more pronounced on *E. coli* than *S. Typhimurium* (Badawy *et al.*, 2004); the Zone Diameter of Inhibition (ZDI) of different honey samples against *E. coli* was 12 to 24 mm and *S. Typhimurium* was 0 to 20 mm. Manuka honey has been demonstrated to be effective against several human pathogens *S. aureus*, *S.*

Typhimurium, *E. coli* and *E. aeruginosa* using agar diffusion method (Lusby *et al.*, 2005; Visavadia *et al.*, 2006). The potency of six varieties of honey from different regions in Algeria were determined against *P. aeruginosa* using disc diffusion method and results showed that Sahara honey have inhibitory activity against tested pathogens and the authors suggested that Sahara honey could be used to manage the wounds and burns infected by *P. aeruginosa* (Boukraa and Niar, 2007). Recently, Boorn *et al.* (2010) tested eleven samples of stingless bee honey and the antibacterial activity was assessed using agar diffusion method which showed inhibitory activity against Gram-negative and Gram-positive bacteria including *S. aureus*, *S. epidermidis*, *S. Typhimurium*, *E. coli* and *P. aeruginosa*. Nilgiris honeys showed Zone Diameter of Inhibition (ZDIs) of 20-21 mm for *S. aureus*, 15-16 mm for *P. aeruginosa* and 13-14 mm for *E. coli* (Rajeswari *et al.*, 2010). RS and Manuka honeys killed *B. subtilis*, *E. coli*, *P. aeruginosa* and *S. aureus* after 24 h of incubation (Kwakman *et al.*, 2011) using disc diffusion method. In this study using disc diffusion method, it was observed that Tualang and Acacia honeys showed higher inhibitory activity compared to Nilgiris honeys, but comparable with the activity from Egyptian clover honey. Tualang and Acacia honeys from Malaysia are able to inhibit the growth of MAR target bacteria using disc diffusion method. Considering the antibacterial activity of honey it was observed that all different honey samples possessed antibacterial activity against target Gram negative and Gram positive pathogenic bacteria as evaluated by disc diffusion and well methods. There were significant differences ($p>0.5$) between target bacteria using disc diffusion method but there was no significant differences ($p<0.5$) with honey type used (Table 2).

The well method indicated that all honey samples significantly ($p<0.5$) inhibited target bacteria, but inhibitory activity varied between bacteria (Table 3 and Fig. 1) with diameter of inhibition between 15.00 and 27.50 mm. The highest inhibitory was shown by H020 from Libya against *S. aureus* (27.50 ± 0.70 mm), *S. Typhimurium* (19.00 ± 1.41 mm), *B. subtilis* (19.50 ± 0.70 mm) and *P. aeruginosa* (18.50 ± 2.12 mm) and from H032 from Malaysia against *E. coli* (17.00 ± 0.00 mm). Zainol *et al.* (2013) tested the antibacterial activity of several Malaysian honeys including Acacia and Tualang and one New Zealand honey (Manuka 18+) using well method against *S. aureus*, *E. coli*, *P.aeruginosa* and *B. cereus*; the results varied from 7.59 to 27.35 mm diameter, the highest inhibitory zone was obtained from Manuka honey against *S. aureus* (19.81 mm) and *E. coli*

(14.04 mm), comparable inhibitory activity from Tualang honey against *P. aeruginosa* (16.22 mm) and *B. cereus* (27.35 mm), while Acacia honey showed the lowest result compared to other samples. In the present study the ability of Al-Seder honey and Acacia honey to kill target bacteria using well method was higher than what observed by Tualang, Acacia and Manuka 18+ as reported by Zainol *et al.* (2013) against all MAR target bacteria. The well method tends to give a higher inhibitory activity for all honey compared to disc diffusion method which may due to the direct attach of the tested compound in liquid form to the agar.

In contrast, when the antibacterial activity of the honey samples were evaluated using Nanophotometer assay no significant difference ($p>0.5$) was observed for both honey and target pathogens. This method used Nanophotometer (IMPLEN) that reported the number of target bacteria cells (both dead and alive) in the sample. All target pathogen reduced in numbers after 24 h incubation compared to control (without honey) (Table 4) ranging from 70 to 100% reduction after 24 h of incubation as shown by Al-Seder honey, 100% (H025), Tualang honey, 100% (H026) and (Acacia honey, 100% (H030) against *S. aureus*.

In order to confirm the inhibitory potency of honey, the growth of target pathogens were evaluated after 24 h incubation in microtiter plates. All honey samples showed antibacterial activity against the target pathogenic bacteria evaluated in which growth was inhibited between 64 to 100% after 24 h of incubation (Table 5). Honey sample Tualang honey (H026) completely inhibited the growth of *S. aureus*, while Acacia honey (H032) showed the highest activity among the evaluated honey samples. Among the pathogens evaluated, *P. aeruginosa* which was more difficult to be inhibited by all the honey samples. It is interesting to note that this test (microtiter plates) give a different results from above (Agar diffusion, well diffusion and nanophotometer methods). Honey samples showed significant differences ($p<0.5$) in antibacterial activity but there was no significant differences ($p>0.5$) between target pathogens in microtiter plates and similar to the results obtained by nanophotometer method.

Other researchers who evaluated antimicrobial activity from sources other than honey, such as extracts from plants or microbial metabolites have included microdilution assay and microtiter plates in addition to the disc and well methods. The disc diffusion method and well method are the two most commonly used to determine antimicrobial properties of honey (Badawy *et al.*, 2004; French *et al.*, 2005;

Lusby *et al.*, 2005; Visavadia *et al.*, 2006; Boukraa and Niar, 2007; Boorn *et al.*, 2010; Rajeswari *et al.*, 2010; Kwakman *et al.*, 2011; Mandal and Mandal, 2011). Based on the results obtained in this study it is suggested that either nanophotometer or microtiter plate assay be included for the evaluation of potency of honey as antibacterial agent.

The above methods (Agar diffusion, well diffusion, nanophotometer and microtiter plate methods) did not evaluate the bactericidal effect of honey; thus plating of survivors of target pathogens after treatment with honey was included by using plate count method (Table 6). Acacia honey H032 and Tualang honey H026 showed total inhibition (survivors <10 est.) against all target photogenic bacteria. *E. coli* was totally inhibited (survivors <10 est.) by all tested honey samples except H031 and H035 which allow recovery of pathogen and reached 10⁵ CFU/ml after 24 h incubation. The potency Tualang honey (H026) and Acacia honey (H032) which showed their ability to kill the Multiple Antibiotic Resistant (MAR) bacteria including *S. aureus*, *B. subtilis*, *S. Typhimurium*, *E. coli* and *P. aeruginosa* suggests their potential to be used as an alternative therapeutic agent in certain medical conditions, particularly wound infection as well as in preservations to control food spoilage.

In most reported study honey was used as it is without heating and at concentrations (w/v) of 15-80% as used by Mavric *et al.* (2008); 50% by Boorn *et al.* (2010), 17.40, 19.20, 20.80, 23.80% by Voidarou *et al.* (2011) and 10, 30, 50, 70, 100% by Moussa *et al.* (2012). The antimicrobial activity was dose-dependent, the higher concentrations the greater the activity affecting both Gram-positive and Gram-negative bacteria. Compounds like glucose oxidase, catalase, ascorbic acid, flavonoids, phenolic acids, organic acids, amino acids and proteins were found present in honey in substantial amounts and can be responsible for the activity of honey (Bogdanov *et al.*, 2008; Perez *et al.*, 2007). However, in this study honey samples were heated at 70°C for 10 min and then filtered using 0.45 µL membrane filters. Heating at 70°C was reported to decrease glucose oxidase but not totally destroyed the enzyme (Kretavicius *et al.*, 2010), whereas hydrogen peroxide activity can be destroyed by heat, light or storage (Bogdanov, 1997). Additionally, all the honey samples were also diluted with deionized water at 200 mg mL⁻¹, a much lower concentration than that used by others and the antibacterial activity was

observed. This study suggested that many compounds including heat stable compounds are responsible for the antibacterial activity and cannot be attributed to one or two main compounds present in honey.

5. CONCLUSION

The antibacterial activity of honey cannot be confirmed by using one single method. While disc diffusion and/or well diffusion method are useful for initial screening, other methods such as microtiter plates, nanophotometer assay and/or plate count method should be included. To ascertain the potency of honey, the survival of target pathogens after treatment with honey should be carried out as well. This study also confirmed that even at low concentration of honey, it still has the potency to inhibit growth of MAR pathogens and this activity could be contributed by water soluble compounds present in honey. Malaysian honeys namely, Tualang and Acacia tend to give the better antibacterial activity compared to other honey samples evaluated. This work further supports that honey could be used as antibacterial agent.

6. ACKNOWLEDGMENT

The researchers would like to thank the management of Faculty of Science and Technology, Islamic Science University of Malaysia for supporting this work.

7. REFERENCES

- Aljadi, A.M. and K.M. Yusoff, 2003. Isolation and identification of phenolic acids in Malaysian honey with antibacterial properties. *Turk. J. Med. Sci.*, 33: 229-236.
- Aween, M.M., Z. Hassan, B.J. Muhiaddin, Y.A. Eljamel and A.S.W. Al-Mabrok *et al.*, 2012a. Antibacterial activity of *Lactobacillus acidophilus* strains isolated from honey marketed in Malaysia against selected Multiple Antibiotic Resistant (MAR) Gram-positive bacteria. *J. Food Sci.*, 77: 364-371. DOI: 10.1111/j.1750-3841.2012.02776.x
- Aween, M.M., Z. Hassan, B.J. Muhiaddin, H.M. Noor and Y.A. Eljamel, 2012b. Evaluation on antibacterial activity of *Lactobacillus acidophilus* strains isolated from honey. *Am. J. Applied Sci.*, 6: 807-817. DOI: 10.3844/ajassp.2012.807.817

- Badawy, O.F.H., S.S.A. Shafii, E.E. Tharwat and A.M. Kamal, 2004. Antibacterial activity of bee honey and its therapeutic usefulness against *Escherichia coli* O157:H7 and *Salmonella typhimurium* infection. *Rev. Scientif. Tech.*, 23: 1011-1122. PMID: 15861897
- Bauer, A.W., W.M. Kirby, J.C. Sherris and M. Turck, 1966. Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.*, 4: 493-496. DOI: 5325707
- Bogdanov, S., 1997. Nature and origin of the antibacterial substances in honey. *LWT Food Sci. Technol.*, 30: 748-753. DOI: 10.1006/fstl.1997.0259
- Bogdanov, S., T. Jurendic, R. Sieber and P. Gallmann, 2008. Honey for nutrition and health: A review. *J. Am. Coll. Nutr.*, 27: 677-689. DOI: 10.1080/07315724.2008.10719745
- Boorn, K.L., Y.Y. Khor, E. Sweetman, F. Tan and T.A. Heard *et al.*, 2010. Antimicrobial activity of honey from the stingless bee *Trigona carbonaria* determined by agar diffusion, agar dilution, broth microdilution and time-kill methodology. *J. Applied Microbiol.*, 108: 1534-1543. DOI: 10.1111/j.1365-2672.2009.04552.x
- Boukraa, L. and A. Niar, 2007. Sahara honey shows higher potency against *Pseudomonas aeruginosa* compared to north Algerian types of honey. *J. Medicinal Food*, 4: 712-4. DOI: 10.1089/jmf.2006.256
- Cooper, R.A., E. Halas, R. Davies, P.C. Molan and K.C. Harding, 2000a. The inhibition of Gram-positive cocci of clinical importance by honey. *Proceedings of the 1st World Healing Congress*, Sept. 10-13, Melbourne, Australia.
- Cooper, R.A., P. Wigley and N.F. Burton, 2000b. Susceptibility of multiresistant strains of *Burkholderia cepacia* to honey. *Lett. Applied Microbiol.*, 31: 20-4. DOI: 10.1046/j.1472-765x.2000.00756.x
- French, V.M., R.A. Cooper and P.C. Molan, 2005. The antibacterial activity of honey against coagulase-negative staphylococci. *J. Antimicrobial Chemotherapy*, 56: 228-231. DOI: 10.1093/jac/dki193
- Hassanain, A.T., A.K. Alyaa and A.J. Karim, 2010. Antimicrobial effect of Malaysian honey on some human pathogens: An *in vitro* study. *Int. Med. J. Malaysia*, 9: 15-17.
- Kretavicius, J., B. Kurtinaitiene, J. Racys and V. Ceksteryte, 2010. Inactivation of glucose oxidase during heat-treatment de-crystallization of honey. *Agriculture*, 97: 115-122.
- Kwakman, P.H.S., A.A. TeVelde, L. de Boer, C.M.J.E. Vandenbroucke-Grauls and S.A.J. Zaat, 2011. Two major medicinal honeys have different mechanisms of bactericidal activity. *Plos One*, 6: 1-7. DOI: 10.1371/journal.pone.0017709
- Lusby, P.E., A.L. Coombes and J.M. Wilkinson, 2005. Bactericidal activity of different honeys against pathogenic bacteria. *Arch. Med. Res.*, 36: 464-467. DOI: 10.1016/j.arcmed.2005.03.038
- Magnusson, J. and J. Schnurer, 2001. *Lactobacillus coryniformis* subsp. *coryniformis* Strain Si3 Produces a broad-spectrum proteinaceous antifungal compound. *Applied Environ. Microbiol.*, 67: 1-5. DOI: 10.1128/AEM.67.1.1-5.2001
- Mandal, M.D. and S. Mandal, 2011. Honey: Its medicinal property and antibacterial activity. *Asian Pacific J. Tropical Biomed.*, 2: 154-160. DOI: 10.1016/S2221-1691(11)60016-6
- Mavric, E., S. Wittmann, G. Barth and T. Henle, 2008. Identification and quantification of methylglyoxal as the dominant antibacterial constituent of Manuka (*Leptospermum scoparium*) honeys from New Zealand. *Molecular Nutr. Food Res.*, 52: 483-9. DOI: 10.1002/mnfr.200700282, PMID: 18210383
- Mohammed, S.J., 2010. Therapeutic efficacy of *Lactobacillus acidophilus* against bacterial isolates from burn wounds. *North Am. J. Medical Sci.*, 2: 1947-2714. PMID: 22558572
- Molan, P.C., 1997. Honey as an Antimicrobial Agent. In: *Bee Products: Properties, Applications and Apitherapy*, Mizrahi, A. and Y. Lensky (Eds.), Plenum Press, New York, pp: 27-37.
- Moussa, A., D., Noureddine, H.S. Mohamed, M. Abdelmelek and A. Saad, 2012. Antibacterial activity of various honey types of Algeria against *Staphylococcus aureus* and *Streptococcus pyogenes*. *Asian Pacific J. Tropical Med.*, 4: 773-776. DOI: 10.1016/S1995-7645(12)60141-2
- Nasir, N.M., A.S., Halim, K.B., Singh, A.A. Dorai and M. M. Haneef, 2010. Antibacterial properties of tualang honey and its effect in burn wound management: A comparative study. *BMC Complementary Alternative Med.*, 10: 31-31. DOI: 10.1186/1472-6882-10-31

- Perez, C., M. Pauli and P. Bazerque, 1990. An antibacterial assay by agar well diffusion method. *Acta Bio Med. Exp.*
- Perez, R.A., M.T., Iglesias, E., Pueyo, M. Gonzalez and C. de Lorenzo, 2007. Amino acid composition and antioxidant capacity of Spanish honeys. *J. Agric. Food Chem.*, 55: 360-365. DOI: 10.1021/jf062055b
- Rahman, M.M., R. Allan and M.S. Azirun, 2010. Antibacterial activity of propolis and honey against *Staphylococcus aureus* and *Escherichia coli*. *Afr. J. Microbiol. Res.*, 4: 1872-1878.
- Rajeswari, T., A. Venugopal, C. Viswanathan, L. Kishmu and C.K. Venil *et al.*, 2010. Antibacterial activity of honey against *Staphylococcus aureus* from infected wounds. *Pharmacologyonline*, 1: 537-541.
- Rozaini, M.Z., A.B.Z. Zuki, M. Noordi, Y. Norimah and A.N. Haki, 2004. The effects of different types of honey on tensile strength evaluation of burn wound tissue healing. *Int. J. Applied Res. Vet. Med.*, 2: 290-296.
- Salvatore, V., S. Christian, C. Francesca, S. Vincenzo and S. Carloand *et al.*, 2010. Antibiotic resistance in *Staphylococcus aureus* and coagulase negative staphylococci isolated from goats with subclinical mastitis. *Vet. Med. Int.* PMID: 20445785
- Subramani, S. and S. Vignesh, 2012. MAR index study and MDR character analysis of a few golden staph isolates. *Asian J. Pharmacy Life Sci.*, 2: 151-154.
- Tan, H.T., R.A. Rahman, S.H. Gan, A.S. Halim and S.A. Hassan *et al.*, 2009. The antibacterial properties of Malaysian tualang honey against wound and enteric microorganisms in comparison to Manuka honey. *BMC Complementary Alternative Med.*, 9: 1-8. DOI: 10.1186/1472-6882-9-34
- Taormina, L.R., B.A. Niemara and L.R. Beuchat, 2001. Inhibitory activity of honey against foodborne pathogens as influenced by the presence of hydrogen peroxide and level of antioxidant power. *Int. J. Food Microbiol.*, 69: 217-225. DOI: 10.1016/S0168-1605(01)00505-0
- Tumin, N.N., A.A. Halimi, M. Shahjahan, N.J. Rani and M.A. Sattar *et al.*, 2005. Antimicrobial activity of local Malaysian honey. *Malaysian J. Pharmaceutical Sci.*, 3: 1-10.
- Visavadia, B.G., J. Honeysett and M.H. Danford, 2006. Manuka honey dressing: An effective treatment for chronic wound infections. *Brazil. J. Maxillofacial Surgery*, 44: 38-41. PMID: 17113690
- Voidarou, C., A. Alexopoulos, S. Plessas, A. Karapanou and I. Mantzourani *et al.*, 2011. Antibacterial activity of different honeys against pathogenic bacteria. *Anaerobe*, 17: 375-379. DOI: 10.1016/j.anaerobe.2011.03.012
- Zainol, M.I., K.M. Yusoff and M.Y.M. Yusof, 2013. Antibacterial activity of selected Malaysian honey. *BMC Complementary Alternative Med.*, 13: 129-129. DOI: 10.1186/1472-6882-13-129